

Attachment to PTO/SB/05 (4/98) Utility Patent Application  
Transmittal

1. METHOD OF REMOVING OFF-FLAVOR FROM FOODS AND DEODORIZER

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
2188  
2189  
2190  
2191  
2192  
2193  
2194  
2195  
2196  
2197  
2198  
2199  
2200  
2201  
2202  
2203  
2204  
2205  
2206  
2207  
2208  
2209  
2210  
2211  
2212  
2213  
2214  
2215  
2216  
2217  
2218  
2219  
2220  
2221  
2222  
2223  
2224  
2225

METHOD OF REMOVING OFF-FLAVOR FROM FOODS  
AND DEODORIZER

Background of the Invention

5       The present invention relates to a method of removing off-flavor from foods such as seafood, meat products and vegetables, a deodorizer, and the like.

      Among the factors undesirably influencing the quality of food are so-called off-flavors, e.g., the fishy smell of seafood,  
10   the fleshy smell of meat, and the grassy smell of soybean. It is known to use sake, wine, shochu, mirin (a sweet sake used as seasoning), etc. in processing or cooking food for the purpose of masking such off-flavors. The deodorizing effect of such substances is considered to be mostly attributable to ethanol.

15       In addition to ethanol mentioned above, the following substances are known to be effective for deodorization: amino acids and organic acids [J. Jap. Soc. for Food Science and Technology, 29(6), 340 (1982)], amino-carbonyl reaction products [J. Fish Sausage, 212, 15 (1982)], saccharides [J. Jap.  
20   Soc. for Food Science and Technology, 46(5), 319 (1999)] and phenol compounds [J. Jap. Soc. for Food Science and Technology, 29(7), 426 (1982)].

      Smoking is known as one of the conventional methods of deodorization utilizing food processing techniques. It is  
25   reported that the above-mentioned phenol compounds are concerned in the deodorization by smoking [Bull. of the Jap. Soc. of Scientific Fisheries, 47(1), 113 (1981); J. Fish Sausage, 212, 15 (1982)].

      Vanillin, which is a kind of phenol compound, is well known  
30   as a substance that overcomes the odor of trimethylamine, off-flavor in seafood. However, the characteristic sweet smell of vanillin is regarded as an undesirable odor in certain kinds of foods and thus prevents the utilization of vanillin as a deodorizer.

35       4-Vinylguaiacol, which is a kind of phenol compound, is a precursor of vanillin. In the field of brewing of sake, wine,

etc., 4-vinylguaiacol itself is considered to be one of the off-flavor components, and studies are made to minimize its formation in the process for producing such brewage.

On the other hand, certain yeast and mold are known as enzyme source of ferulic acid decarboxylase having the activity to form 4-vinylguaiacol from ferulic acid, and the ferulic acid decarboxylase gene of yeast has been obtained [Gene, 142, 107 (1994); Japanese Published Unexamined Patent Application No. 276788/98].

#### Summary of the Invention

Some of the off-flavors of foods are inherent in food materials, and others are developed in the course of processing or cooking foods. A known example of the latter is the one developed by oxidation of lipid caused by heating. As the off-flavor developed by oxidation can be reduced by suppressing oxidation, it is considered as one of the effective methods for removing off-flavor to cause a substance having antioxidant activity to be present in the system.

An object of the present invention is to provide an effective method of removing off-flavor from foods such as seafood, meat products and vegetables, and to provide deodorizers, etc.

Another object of the present invention is to provide products containing deodorizers of the present invention.

One of other object of the present invention is to provide a method of producing a deodorizer comprising an oxidation reaction step for conducting an oxidation reaction of a polymer of phenol compounds having a styrene structure in the presence of oxygen, and a deodorizer produced by the method thereof.

One of other object of the present invention is to provide a method of producing a seasoning comprising an oxidation reaction step for conducting an oxidation reaction of a polymer of phenol compounds having a styrene structure in the presence of oxygen, and a seasoning produced by the method thereof.

One of other object of the present invention is to provide

a method of producing liquor comprising an oxidation reaction step for conducting an oxidation reaction of the resultant liquid of the separation step in the presence of oxygen, and liquor produced by the method thereof.

5 In one embodiment, the present invention provides a method of removing off-flavor from a food which comprises causing a polymer of phenol compounds having a styrene structure to be present in the food.

10 In another embodiment, the invention provides a deodorizer comprising a polymer of phenol compounds having a styrene structure.

In another embodiment, the invention provides a seasoning comprising a polymer of phenol compounds having a styrene structure.

15 In another embodiment, the invention provides a cosmetic comprising a polymer of phenol compounds having a styrene structure.

20 In another embodiment, the invention provides an antioxidant comprising a polymer of phenol compounds having a styrene structure.

In another embodiment, the invention provides a method of producing a polymer of phenol compounds having a styrene structure, which comprises converting phenol compounds having a styrene structure into the polymer in the presence of oxygen.

25 In another embodiment, the invention provides a method of producing a seasoning comprising a polymer of phenol compounds having a styrene structure, which comprises causing phenol compounds having a styrene structure to be present in any of the steps in a process for producing a seasoning, and  
30 conducting an oxidation reaction of the phenol compounds in the presence of oxygen during or after the completion of the process.

In another embodiment, the invention provides a method of producing a deodorized food, which comprises cooking a food using the above deodorizer.


35 In another embodiment, the invention provides a method of deodorizing a food, which comprises cooking the food using

the above deodorizer.

In another embodiment, the invention provides a method of producing liquor, which comprises an oxidation reaction step for conducting an oxidation reaction of the resultant liquid of the separation step in the presence of oxygen.



#### Detailed Description of the Invention



The phenol compounds having a styrene structure may be any compounds having hydroxy and vinyl on a benzene ring (said vinyl may be substituted by carboxy or the like). Preferred are compounds represented by general formula (I):





(wherein  $R^1$  represents hydrogen or carboxy, and  $R^2$  represents hydrogen or lower alkoxy). The lower alkyl moiety of the above lower alkoxy includes straight-chain or branched alkyl having 1 to 6 carbon atoms, such as methyl, ethyl, propyl, isopropyl, butyl, pentyl and hexyl. In the polymer of compounds represented by general formula (I), the constituent compounds may be the same or different.

The polymers of phenol compounds having a styrene structure include dimers, trimers, tetramers and pentamers of phenol compounds having a styrene structure. Preferred are dimers. More preferred are those represented by the following formulae (II) to (VII):







(wherein  $R^{1a}$ ,  $R^{1b}$ ,  $R^{1c}$ ,  $R^{1d}$ ,  $R^{1e}$ ,  $R^{1f}$ ,  $R^{1g}$ ,  $R^{1h}$ ,  $R^{1i}$ ,  $R^{1j}$ ,  $R^{1k}$  and  $R^{1l}$ , which may be the same or different, each represents hydrogen or carboxy, and  $R^{2a}$ ,  $R^{2b}$ ,  $R^{2c}$ ,  $R^{2d}$ ,  $R^{2e}$ ,  $R^{2f}$ ,  $R^{2g}$ ,  $R^{2h}$ ,  $R^{2i}$ ,  $R^{2j}$ ,  $R^{2k}$  and  $R^{2l}$ , which may be the same or different, each represents hydrogen or lower alkoxy). Particularly preferred is 2,3-dihydro-2-(4-hydroxy-3-methoxyphenyl)-7-methoxy-5-vinylbenzofuran (VGD).



### Method of Obtaining Phenol Compounds Having a Styrene Structure

The phenol compounds having a styrene structure can be synthesized according to known methods. For example, 4-vinylphenol can be obtained by reduction, dehydration and hydrolysis of 4-acetoxyacetophenone. Ferulic acid can be obtained by subjecting 4-hydroxy-3-methoxybenzaldehyde and malonic acid to condensation decarboxylation in pyridine in the presence of piperidine. p-Coumaric acid can be obtained from p-aminocinnamic acid via diazonium salt, or from p-oxybenzaldehyde by the Perkin reaction. 4-Vinylguaiacol can be obtained by preparing 4-acetoxy-3-methoxycinnamic acid from vanillin using acetic anhydride or sodium acetate, followed by hydrolysis and decarboxylation using quinoline.

Some phenol compounds having a styrene structure, e.g. ferulic acid, are also commercially available.

4-Vinylguaiacol, ferulic acid, etc. can also be produced from natural products including food materials.

The methods of producing these phenol compounds from natural products are described below.

### Method of Producing Ferulic Acid

Ferulic acid can be obtained by decomposing ferulic acid ester existing in fibrous substances such as arabinoxylan and pectin, which are one of the cell wall components of plants, in the presence of ferulic acid esterase in the usual way. Examples of plants known to contain ferulic acid or ferulic acid ester are cereals (e.g. rice, wheat, barley and buckwheat), potatoes (e.g. potatoes and sweet potatoes), fruits (e.g. grapes, apples and citrus fruits) and vegetables.

Ferulic acid can also be obtained by treating the above plants with fibrous hydrolases such as xylanase, arabinase, cellulase, pectinase and hemicellulase in the usual way.

Further, ferulic acid can be obtained, for example, by decomposing lignin in wood by ordinary methods such as thermal decomposition, ethanol decomposition and use of hydrolases (e.g.

ligninase, manganese peroxidase and laccase).

#### Method of Producing 4-Vinylguaiacol

4-Vinylguaiacol can be obtained by treating ferulic acid  
5 in the presence of ferulic acid decarboxylase in the usual way.

Useful enzyme sources of ferulic acid decarboxylase  
include not only the enzyme itself, but also cells of filamentous  
fungi, yeast, etc. having the enzyme activity and crude enzyme  
extracted from such cells by appropriate means [Appl. Environ.  
10 Microbiol., 59(7), 2244 (1993)]. Preferred yeast strains are  
those having the enzyme activity of 0.5 unit/g (unit is defined  
in the explanation of Examples hereinbelow) (wet cell weight)  
or more. Examples of such yeast strains are Saccharomyces  
cerevisiae IFO 2260, IFO 1953 and IFO 0233 and Dia Yeast  
15 (trademark for pressed yeast produced by Kyowa Hakko Kogyo Co.,  
Ltd.).

#### (1) Method of Producing the Polymer of Phenol Compounds Having a Styrene Structure

20 The method of producing the polymer of phenol compounds  
having a styrene structure is described below.

The conversion of phenol compounds having styrene  
structure into the polymer is carried out, for example, by  
maturing at 10 to 80°C in a solvent (e.g. a solvent containing  
25 ethanol) in the presence of oxygen. For instance, the  
conversion of 5 ppm 4-vinylguaiacol into VGD is preferably  
carried out by maturing at 25°C for 6 months or more, at 37°C  
for 4 months or more, or at 60°C for 20 days or more.

30 It is preferred to make aeration or blow oxygen into the  
reaction system in view of reduction in reaction time.

The above conversion reaction can be accelerated under  
irradiation with light, and further accelerated in the presence  
of a photosensitizer.

35 When the reaction is carried out under irradiation with  
light in the presence of a photosensitizer, the intensity and  
time of irradiation suitable for the conversion at 5 to 80°C,

preferably 10 to 30°C, are selected. For instance, the conversion of 1 liter of a solution containing 5 ppm 4-vinylguaiacol into VGD is preferably carried out by treatment at 20°C under a fluorescent light of 20 to 40 W for 1 to 3 days, or at 30°C under a fluorescent light of 20 to 40 W for 0.5 to 2 days.

Preferred light sources are those mainly emitting light with wavelengths in the range of 240 to 500 nm. Examples of such light sources are sunlight, a fluorescent light, an incandescent lamp, a black light lamp, a sterilization lamp and an insect-drawing lamp, which may be used alone or in combination. Particularly, a black light lamp, a sterilization lamp and an insect-drawing lamp are preferred in respect of energy efficiency.

Hydroxymethylfurfural and riboflavin are able to shown as an example of photosensitizer. The conversion rate varies according to the combination of photosensitizer and light source. Preferred combinations are ultraviolet light and hydroxymethylfurfural, riboflavin, etc., and visible light and riboflavin, etc. If a fermentation liquor to be irradiated with light contains riboflavin, whether it is contained in raw materials or formed by fermentation, it is not necessary to add a photosensitizer. The concentration of the photosensitizer is preferably 1 ppb or more, more preferably 10 to 5000 ppb.

The above conversion reaction can also be carried out in the presence of an enzyme capable of converting the phenol compounds having a styrene structure into the polymer.

Useful enzyme sources of such enzymes (e.g. peroxidase) include a peroxidase reagent produced by ICN Biomedicals Inc. and a crude enzyme solution of peroxidase extracted from horseradish with water.

Recovery of the polymer (hereinafter the polymer of phenol compounds having a styrene structure is sometimes referred to merely as the polymer) from the reaction mixture is carried out, for example, by using a nonionic porous resin such as Diaion HP 20 (Mitsubishi Chemical Corporation) (the

"nonionic porous resin" as used herein means a resin which does not have a functional group such as an ion-exchange group and adsorbs various organic substances with the van der Waals force). The nonionic porous resin is charged into a column and is  
5 equilibrated with 5 to 30% aqueous ethanol or methanol. The reaction mixture is passed through the column to adsorb the polymer on the resin, followed by elution of the polymer with alcohol. If necessary, the elute is dried under reduced pressure to obtain the desired product as a powder.

10

### (2) Method of Removing Off-flavor from Foods

In accordance with the present invention, off-flavor in foods can be reduced or completely removed by causing the polymer of phenol compounds having a styrene structure and /or the  
15 reaction mixture comprising the polymer of phenol compounds having a styrene structure obtained by the above method to be present in the foods, for example, by addition of the polymer to the foods.

In this method, not only the polymer in a pure form but  
20 also a substance containing the polymer can be employed. For example, the use of the reaction mixture described above and also a crude purified form of the reaction mixture can be employed. For the use of the deodorizer or seasoning of the present invention described below in removal of off-flavor from  
25 foods is included within the scope of the method of removing off-flavor from foods.

In the method of removing off-flavor from foods according to the present invention, it is preferred to cause the polymer to be present in the foods at a concentration of 0.5 ppb or more.

30

### (3) Deodorizer

The deodorizer of the present invention is effective for the deodorization of off-flavors in foods such as the fishy smell of seafood, the fleshy smell of meat and the grassy smell of  
35 soybean, body odors such as odor caused by maturation and the smell of armpit, etc.

The deodorizer of the present invention can be obtained by mixing the polymer of phenol compounds having a styrene structure with an appropriate solvent (e.g. water, ethanol, and a mixture thereof) or powder (an excipient for the use for foods, e.g. starch, dextrin and lactose, or an excipient for the use as cosmetics, e.g. talc). The solution of the polymer in ethanol or the like, which is obtained in the step of purification from the reaction mixture in the process for producing the polymer, and/or the reaction mixture itself, as such, can also be used as the deodorizer of the present invention. The polymer of phenol compounds having a styrene structure can be obtained by the process described in (1) above or processes similar thereto.

The deodorizer of the present invention may comprise various ingredients according to the use and purpose, in addition to the polymer of phenol compounds having a styrene structure. When the deodorizer is used for the deodorization of foods, it may be, for example, added to foods in the form of a mixture with ethanol, or used as the seasoning of the present invention described below.

When applied to foods, the deodorizer of the present invention may be added in the form of a solution containing the polymer in 1 to 80%, preferably 5 to 30% aqueous ethanol, preferably at a concentration of 10 ppb or more, more preferably 10 ppb to 100 ppm.

For the purpose of removing body odor, the deodorizer of the present invention can be used, for example, as the cosmetic of the present invention described below.

#### (4) Seasoning

The seasonings of the present invention can be obtained by causing the polymer of phenol compounds having a styrene structure to be present in known seasonings currently used or novel seasonings which will be developed in the future preferably at a concentration of 10 ppb or more, more preferably 10 ppb to 100 ppm. Specifically, the seasonings of the present

invention can be obtained by adding the polymer of phenol compounds having a styrene structure to a seasoning as a final product or to materials in any of the steps in a process for producing a seasoning, or by causing phenol compounds having a styrene structure to be present in any of the steps in a process for producing a seasoning, and conducting an oxidation reaction of the phenol compounds in the presence of oxygen during or after the completion of the process. The polymer of phenol compounds having a styrene structure can be obtained by the process described in (1) above or processes similar thereto. The process for conducting an oxidation reaction of the phenol compounds in the presence of oxygen during or after the completion of the process is described in detail in the description of fermented seasoning below.

The preferred seasonings according to the present invention are those containing at least one member of amino acids (e.g. glutamic acid, aspartic acid, alanine, arginine, leucine, tyrosine and glycine), organic acids (e.g. succinic acid, lactic acid, malic acid, citric acid and acetic acid), alcohols (e.g. ethanol, 1-propanol, 2-methylpropanol and isoamyl alcohol), saccharides (e.g. glucose, fructose, sucrose and maltose) and esters (e.g. isoamyl acetate and ethyl caproate). The seasonings of the present invention may be in the form of liquors (e.g. sake, shochu, whiskey and brandy), fermented seasonings, extract seasonings, etc.

#### Extract Seasoning

The extract seasonings of the present invention are seasonings containing stocks obtained by boiling or smothering meat, marine products, vegetables, etc. or concentrates of the stocks among the seasonings of the present invention described above. Such seasonings include meat extract seasonings, marine product extract seasonings and vegetable extract seasonings. The meat extract seasonings are produced mainly from meat or bones of cattle, pigs, fowls, etc., the marine product extract seasonings are produced mainly from marine products (e.g.

bonitos, mackerels, tuna, sardines, croakers, pike congers, thread sea bream, scallops, short-necked clams, oysters, krills and crabs) or dried products thereof, and vegetable extract seasonings are produced mainly from vegetables (e.g. onion, garlic, carrots, Chinese cabbages, cabbages, *shiitake* mushrooms and tangleweed) or dried products thereof, according to conventional methods.

The extract seasonings of the present invention can be produced in the following manner. An extract material (e.g. dried bonito) is subjected to extraction with a solution containing the polymer of phenol compounds having a styrene structure in ethanol, water, or a mixture thereof (a preferred example is the cooking liquor of the present invention described below) at 5 to 120°C, preferably 20 to 80°C for 1 to 10 hours, followed by solid-liquid separation to obtain the supernatant as an extract. The obtained extract is concentrated by heating or freezing, by use of membrane, or under reduced pressure, as may be required, whereby the extract seasonings of the present invention are obtained as concentrates.

#### Fermented Seasoning

The fermented seasonings of the present invention are seasonings produced by utilizing the fermentation by microorganisms or the like among the seasonings of the present invention described above. These seasonings may take various forms such as vinegar, cooking liquor, *sake*, wine, *mirin*, soy sauce and *miso*.

The fermented seasonings of the present invention can be obtained by adding the polymer of phenol compounds having a styrene structure to a fermented seasoning as a final product or to materials in any of the steps in a process for producing a fermented seasoning. However, they are preferably produced by conducting an oxidation reaction of the phenol compounds in the presence of oxygen during or after the completion of the production process for producing a fermented seasoning.

Production of the fermented seasonings of the present

invention comprising conducting an oxidation reaction of the phenol compounds in the presence of oxygen during or after the completion of the process for producing a fermented seasoning is exemplified by a process which comprises forming 4-vinylguaiacol during the production process, and after the completion of the production process, converting 4-vinylguaiacol in the obtained fermented seasoning into VGD. This process is explained below referring to the production of a cooking liquor.

As the first step, cereals such as rice (usually containing ferulic acid or ester thereof) as a raw material are liquefied and saccharified using a filamentous fungus such as rice *koji*. In this step, liquefying enzyme such as  $\alpha$ -amylase and saccharifying enzyme such as glucoamylase may be added if necessary. Then, yeast is added to cause fermentation and the obtained fermentation liquor (*moromi*) is filtered to obtain a cooking liquor as the filtrate. The filamentous fungus used above has also fibrous hydrolase activity and ferulic acid esterase activity and contributes to decomposition of fibers in cereals and liberation of ferulic acid.

In the above step, ferulic acid is formed from cereals in the presence of xylanase together with glucoamylase. The formed ferulic acid is converted into 4-vinylguaiacol by ferulic acid decarboxylase contained in specific yeasts such as those mentioned above in the method of producing 4-vinylguaiacol. The resultant 4-vinylguaiacol is contained in the produced cooking liquor. Filamentous fungi is also utilized in this step in place of yeast fungi in order to use the same enzyme activity, ferulic acid decarboxylase. When ferulic acid decarboxylase contained in filamentous fungi is used, the addition of yeast mentioned above is not necessary.

As the next step, the cooking liquor containing 4-vinylguaiacol is subjected to reaction for converting 4-vinylguaiacol into a polymer such as VGD, whereby the cooking liquor of the present invention containing VGD is obtained. The reaction for converting 4-vinylguaiacol into the polymer can



be carried out by the process described in (1) above or processes similar thereto.

The polymer may also be formed during the process for producing a fermented seasoning by adding or forming the phenol compounds having a styrene structure at an appropriate time during the production process and carrying out reaction for converting the phenol compounds into the polymer in the same manner as above at an appropriate time during the production process.

#### (5) Cosmetic

The cosmetics of the present invention are formulated to contain the polymer of phenol compounds having a styrene structure preferably at a concentration of 10 ppb or more, more preferably 10 ppb to 1 ppm, based on the total weight of the product. The polymer of phenol compounds having a styrene structure and/or the reaction mixture containing the polymer of phenol compounds having a styrene structure can be obtained by the process for producing the polymer described above or processes similar thereto.

The cosmetics of the present invention may further comprise cosmetic bases such as pigments, perfumes, preservatives, surfactants, antioxidants and ultraviolet absorbents, as may be required. The cosmetics can be prepared, for example, according to the methods described in "The development manual of preparations for percutaneous application " 1st ed., Mitsuo Matsumoto ed., [Seishi Shoin (1985)].

Examples of the pigments include tar pigments, iron oxide, titanium oxide and zinc oxide. Examples of the perfumes include animal perfumes such as musk, plant perfumes such as peppermint oil, lemon oil and rose oil, and synthetic perfumes such as benzyl alcohol and anisole. Examples of the preservatives include paraben, methylparaben, ethyl p-oxybenzoate and butyl p-oxybenzoate. Examples of the surfactants include anionic surfactants such as sodium cetyl sulfate, nonionic surfactants

such as polyoxyethylene alkyl ether, polyoxyethylene fatty acid ester, polyoxyethylene polyhydric alcohol fatty acid ester, fatty acid ester hardened castor oil, polyhydric alcohol fatty acid ester and polyglycerin fatty acid ester, cationic surfactants such as tetraalkylammonium salt, and amphoteric surfactants such as betaine surfactants, sulfobetaine surfactants, sulfoamino acid surfactants and sodium N-stearoyl-L-glutamate. An example of the antioxidant is dibutylhydroxytoluene. Examples of the ultraviolet absorbents include 2-ethylhexyl p-methoxycinnamate and 4-tert-butyl-4'-methoxydibenzoylmethane.

The cosmetics may take various forms such as cream, emulsion, lotion, essence, enriched lotion and pack. In the case of emulsion, for example, the cosmetic can be produced by a conventional method, that is, by emulsifying an oil phase and a water phase respectively heated and then cooling the resultant emulsion.

The cosmetics of the present invention are effective for the removal of body odor, etc.

#### (6) Antioxidant

The antioxidant of the present invention contains the polymer of phenol compounds having a styrene structure preferably at a concentration of 10 ppb or more, more preferably 10 ppb to 100 ppm.

The antioxidant of the present invention can be obtained by mixing the polymer of phenol compounds having a styrene structure with an appropriate solvent (e.g. water, ethanol, and a mixture thereof) or powder [e.g. those mentioned in the explanation of powder in (3)]. The solution of the polymer in ethanol or the like which is obtained in the step of purification from the reaction mixture in the process for producing the polymer, as such, can also be used as the antioxidant of the present invention. The polymer of phenol compounds having a styrene structure can be obtained by the method of producing the polymer described above or processes similar thereto.

The antioxidant of the present invention can be used, for example, by adding it to foods, seasonings, cosmetics, etc. as final products or to materials during the processes for producing them, or by adding it to foods during cooking.

5 The antioxidant of the present invention is useful in preventing oxidation of foods, seasonings, cosmetics, etc. For example, it is useful in preventing oxidation of linoleic acid in foods.

10 (7) Method of Producing a Deodorized Food Which Comprises Cooking a Food Using the Above Deodorizer and a Food Obtained by this Method

The method of producing a deodorized food which comprises cooking a food using the above deodorizer can be carried out, for example, by using the deodorizer described in (3) above in cooking seafood, meat or vegetables (e.g. soybean) preferably in such an amount that the concentration of the polymer of phenol compounds having a styrene structure in the food becomes 0.5 ppb or more. The foods of the present invention which can be  
15 obtained by this method are desirable in that they are freed of off-flavor.

20 (8) Method of Deodorizing a Food Which Comprises Cooking the Food Using the Above Deodorizer

25 The method of deodorizing a food which comprises cooking the food using the above deodorizer can be carried out, for example, by using the deodorizer described in (3) above in cooking seafood, meat or vegetables (e.g. soybean) preferably in such an amount that the concentration of the polymer of phenol  
30 compounds having a styrene structure in the food becomes 0.5 ppb or more.

Certain embodiments of the present invention are illustrated in the following examples.

35 In the following examples, high performance liquid chromatography (HPLC) was carried out under the following

conditions.

Column: Inertsil ODS - 24.6 x 250 mm (GL Sciences)

Detector: OD 280 nm

- 5           The definitions of enzyme activities as used in the following examples are as follows.

#### Ferulic Acid Decarboxylase Activity

10           To 0.9 ml of a culture of test yeast obtained by culturing the yeast in YPD medium at 30°C overnight with shaking was added 0.1 ml of an aqueous solution containing 1 g/liter ferulic acid (Ichimaru Falcos), and the mixture was allowed to stand at 25°C. The amount of 4-vinylguaiacol in the supernatant of the resultant reaction mixture was determined by HPLC. One unit  
15           of the enzyme activity was defined as that amount of the enzyme which forms 1 ppm of 4-vinylguaiacol per hour.

#### Xylanase Activity

20           Xylan derived from oats (100 mg, Nacalai Chemicals) was dissolved in 9.0 ml of 50 mol/liter acetate buffer (pH 4.5), followed by addition of 1.0 ml of a test enzyme solution. The mixture was subjected to reaction at 52°C with stirring, and the change in reducing sugar equivalent in the reaction mixture was determined by the dinitrosalicylic acid reaction. One unit  
25           of the enzyme activity was defined as that amount of the enzyme which liberates 1  $\mu$ mol of reducing sugar equivalent from xylan in one minute.

#### Determination of the Phenol Compounds

30           In the following examples, analysis of the phenol compounds having a styrene structure was carried out in the following manner.

35           A test substance was analyzed by HPLC in the following manner. In the analysis of VGD, a sample was concentrated to make a test solution. The sample (50 ml) was passed through a C<sub>18</sub> cartridge for HPLC pretreatment and eluted with 0.8 ml

of acetonitrile. The eluate was brought to a volume of 1.0 ml and then analyzed under the above conditions.

#### Fractionation Conditions in HPLC

5 Mobile phase: solution A: 20 mmol/liter aqueous solution of sodium acetate

solution B: acetonitrile

Elution was carried out with linear gradient of solution B/solution A (volume ratio) of 30 to 80%/25 minutes.

Column temperature: 40°C

Flow rate: 1.0 ml/minute

#### Seven Points Scoring and Test of Significance

15 In the following examples, the term "seven points scoring" refers to a method of evaluating a subject with respect to the strength of fishy smell by giving the subject scores of one point (weak) to seven points (strong). Student t-test was made as to the significance of difference. The symbol n refers to the number of persons.

#### Example 1

Eight hundred milliliters of 7.75% aqueous ethanol containing 50 ppm 4-vinylguaiacol (4-VG) and 41.6 ppm hydroxymethylfurfural was irradiated with ultraviolet light for 40 minutes. The resultant reaction mixture was passed through Diaion HP 20 (trademark, Mitsubishi Chemical Corporation) equilibrated with 7.75% aqueous ethanol, followed by washing with 80% aqueous methanol. A fraction eluted with 8 ml of methanol (fraction A) was obtained as a crude product.

Separately, 20 liters of 7.75% aqueous ethanol containing 50 ppm 4-vinylguaiacol and 0.5 ppm riboflavin was kept in a thermostat with fluorescent illumination at a reaction temperature of 25°C for 24 hours. The resultant reaction mixture was passed through Diaion HP 20 (trademark, Mitsubishi Chemical Corporation) equilibrated with 7.75% aqueous ethanol,

followed by washing with 80% aqueous methanol. A fraction eluted with 200 ml of methanol (fraction B) was obtained as a crude product.

The fractions obtained above were concentrated 10-fold under reduced pressure and then fractionated by C<sub>18</sub> HPLC, whereby a purified VGD fraction (fraction C) was obtained for each of the above fractions. Identification of the obtained VGD was carried out.

The fractionation conditions in HPLC and the result of identification of VGD are shown below.

#### Fractionation Conditions in HPLC

Mobile phase: 20 mmol/liter aqueous solution of sodium acetate/acetonitrile = 45/55

Column temperature: 40°C

Flow rate: 1.0 ml/minute

#### Physical Properties of VGD

##### FAB mass spectrum:

Positive mode: m/z 299 (M+H)<sup>+</sup>

Negative mode: m/z 297 (M-H)<sup>-</sup>

<sup>1</sup>H-NMR (400 MHz, CD<sub>3</sub>OD); [δ ppm (integration, multiplicity, coupling constant J (Hz))]; 6.98 (1H, d, 2.0), 6.92 (1H, br.s), 6.91 (1H, br.s), 6.84 (1H, dd, 8.1, 2.0), 6.78 (1H, d, 8.1), 6.64 (1H, dd, 17.6, 11.0), 5.69 (1H, t, 8.9), 5.60 (1H, dd, 17.6, 1.1), 5.06 (1H, dd, 11.0, 1.1), 3.86 (3H, s), 3.82 (3H, s), 3.55 (1H, dd, 15.6, 9.3), 3.16 (1H, dd, 15.6, 8.5)

<sup>13</sup>C-NMR (100 MHz, CD<sub>3</sub>OD); δ ppm (multiplicity); 149.1 (s), 149.1 (s), 147.7 (s), 145.4 (s), 138.1 (d), 134.5 (s), 133.2 (s), 129.7 (s), 120.0 (d), 116.4 (d), 116.2 (d), 111.4 (d), 111.4 (t), 110.7 (d), 86.7 (d), 56.7 (q), 56.4 (q), 39.4 (t)

#### Example 2

Twenty liters of 7.75% aqueous ethanol containing 50 ppm 4-vinylguaiacol and 0.5 ppm riboflavin was kept in a thermostat with fluorescent illumination at a reaction temperature of 25°C

for 24 hours. The resultant reaction mixture was passed through Diaion HP 20 (trademark, Mitsubishi Chemical Corporation) equilibrated with 7.75% aqueous ethanol. After washing with 80% aqueous ethanol, elution was carried out with 200 ml of ethanol to obtain a deodorizer of the present invention. The VGD concentration in this deodorizer was 40.4 ppm.

### Example 3

Two mixtures were prepared, each consisting of 100 g of onion, 15 g of carrot, 15 g of Welsh onion, 10 g of parsley, 0.5 g of whole pepper, a laurel leaf, 2.5 g of salt, 700 ml of water and 250 g of bony parts of sea bream. To one of the mixtures was added 100 ml of the deodorizer of the present invention obtained in Example 2 (VGD concentration: 40.4 ppm) and to the other (control) was added 100 ml of ethanol. Each of the mixtures was boiled for 20 minutes and then filtered through cloth. The obtained soup was kept at 60°C and evaluated by seven points scoring with respect to the strength of fishy smell perceived when the lid of the vessel was taken off. The results are shown in Table 1. The soup prepared with addition of VGD of the present invention had a significantly reduced fishy smell, compared with the control.

Table 1

	VGD-treated soup	Control
Average score	2.75*	4.00
Standard deviation	0.46	0.76

\* n=8      \*: P<0.01

### Example 4

To 2.0 liters of water were added 370 g of rice *koji*, 1.5 g of yeast (Dia Yeast; trademark, Kyowa Hakko Kogyo Co., Ltd.) and 7 g of lactic acid, followed by fermentation at 20°C for 2 days to obtain a yeast culture (first stage fermentation). To 1.3 liter of water were added 660 g of rice and 2 g of Spitase CP-40G (trademark for  $\alpha$ -amylase preparation produced by Nagase Seikagaku Kogyo Co., Ltd., 400000 units/g), and the mixture was

liquefied by heating at 90°C for 30 minutes and then cooled to 50°C. To the resultant mixture were added 2 g of Sumizyme 3000 (trademark for glucoamylase produced by Shinnihon Kagaku Co., Ltd., 3000 units/g) and 2 g of Sumizyme AC (trademark for hemicellulase produced by Shinnihon Kagaku Co., Ltd., 6400 units/g), followed by reaction at 50°C for 16 hours to obtain a saccharified rice liquor. This saccharified rice liquor and 370 g of rice *koji* were added to the yeast culture, followed by fermentation at 20°C for 2 days (second stage fermentation).

To 4.4 liters of water were added 1800 g of rice and 7 g of Spitase CP-40G, and the mixture was liquefied by heating at 90°C for 30 minutes and then cooled to 50°C. To the resultant mixture were added 7 g of Sumizyme 3000 and 7 g of Sumizyme AC, followed by reaction at 50°C for 16 hours to obtain a saccharified rice liquor. This saccharified rice liquor was added to the fermentation liquor obtained by the second stage fermentation, followed by fermentation at 20°C for 7 days (third stage fermentation). The obtained fermentation product was filtered and the filtrate (10 liter) was pasteurized (hereinafter the filtrate prepared in this manner is referred to as *moromi*).

The amount of 4-vinylguaiacol formed in the fermentation process was 2.1 ppm at the start of the third stage fermentation and 5.1 ppm after 7 days of fermentation.

The *moromi* obtained above (containing 5 ppm 4-vinylguaiacol) was kept in a thermostat with fluorescent illumination at 25°C for 24 hours to obtain a cooking liquor of the present invention. The cooking liquor contained 0.1 ppm 4-vinylguaiacol and 67 ppb VGD.

### Example 5

To 1 liter of the *moromi* prepared in Example 4 was added 1.2 g of a peroxidase reagent (81.8 units/mg, ICN Biomedicals Inc.), followed by reaction at 25°C for 48 hours to obtain a cooking liquor of the present invention. The cooking liquor contained 0.2 ppm 4-vinylguaiacol and 66 ppb VGD.



Example 6

Commercial materials for oden (Japanese hotchpotch) with soup were cooked using the cooking liquor of the present invention obtained in Example 4 in an amount of 10 wt% based on the soup. The cooked oden was evaluated by seven points scoring with respect to the strength of fishy smell perceived when the lid of the pot was taken off. As a control, the same procedure as above was repeated except for the use of a commercial cooking liquor instead of the cooking liquor of the present invention. The results are shown in Table 2. The cooking liquor of the present invention significantly reduced fishy smell, compared with the commercial one.

Table 2

	Cooking liquor of the invention	Commercial cooking liquor
Average score	3.00*	4.50
Standard deviation	1.30	0.70

\* n=10

\*: P&lt;0.05

Example 7

Two mixtures were prepared, each consisting of 100 g of onion, 15 g of carrot, 15 g of Welsh onion, 10 g of parsley, 0.5 g of whole pepper, a laurel leaf, 2.5 g of salt, 700 ml of water and 250 g of bony parts of sea bream. To one of the mixtures was added 100 ml of the cooking liquor of the present invention obtained in Example 4 and to the other was added 100 ml of a commercial cooking liquor. Each of the mixtures was boiled for 20 minutes and then filtered through cloth. The obtained soup was kept in a vessel at 60°C and evaluated by seven points scoring with respect to the strength of fishy smell perceived when the lid of the vessel was taken off. The results are shown in Table 3. The cooking liquor of the present invention significantly reduced fishy smell, compared with the commercial one.

Table 3

	Cooking liquor of the invention	Commercial cooking liquor
Average score	3.47*	4.93
Standard deviation	1.06	1.16

\* n=15      \*: P<0.01

#### Example 8

To 350 g of well washed short-necked clams with shells was added 50 ml of the cooking liquor of the present invention obtained in Example 4. The short-necked clams were brought to the boil on a high flame and then heated on a low flame for one minute. The thus cooked short-necked clams were kept at 60°C and evaluated by seven points scoring with respect to the strength of fishy smell perceived when the lid of the vessel was taken off. The same procedure as above was repeated except for the use of a commercial cooking liquor instead of the cooking liquor of the present invention. The results are shown in Table 4. The cooking liquor of the present invention significantly reduced fishy smell, compared with the commercial one.

Table 4

	Cooking liquor of the invention	Commercial cooking liquor
Average score	3.50*	5.13
Standard deviation	1.41	1.13

\* n=7      \*: P<0.05

#### Example 9

A mixture of 5 kg of dried bonito, 1 liter of the cooking liquor of the present invention obtained in Example 4, 1.4 liter of 59% denatured alcohol and 7.6 liter of water was kept at 40°C for 2 hours, followed by solid-liquid separation to obtain bonito extract. The obtained bonito extract was evaluated by seven points scoring with respect to the strength of fishy smell while being kept at 60°C. The same procedure as above was repeated except for the use of a commercial cooking liquor instead of the cooking liquor of the present invention. The results are shown in Table 5. The cooking liquor of the present

invention significantly reduced fishy smell, compared with the commercial one.

Table 5

	Cooking liquor of the invention	Commercial cooking liquor
Average score	3.47*	4.27
Standard deviation	1.46	0.80

5 \* n=15      \*: P<0.05

#### Example 10

A mixture was prepared, which consisted of 0.2 g of methyl p-oxybenzoate, 3.0 g of 1,3-butylene glycol, 0.8 g of sodium citrate, 0.1 g of ethylenediaminetetraacetic acid (EDTA) disodium salt, 0.4 g of polyoxyethylene hardened castor oil, 5.0 g of denatured alcohol, 0.05 g of perfume and 0.5 ml of the deodorizer of the present invention obtained in Example 2. The mixture was made up to 100 ml with purified water to obtain a lotion of the present invention. Ten male adults were made the subjects of this experiment. The obtained lotion (1.0 ml) was infiltrated into absorbent cotton and applied to the armpit of each subject, followed by evaluation of the odor of armpit by seven points scoring. The results are shown in Table 6. The lotion of the present invention significantly reduced unpleasant odor of armpit, compared with the control lotion containing no VGD.

Table 6

	Lotion of the invention	Commercial lotion
Average score	3.60*	4.80
Standard deviation	1.20	1.40

25 \* n=10      \*: P<0.05

#### Example 11

Antioxidation test was carried out on an antioxidant of the present invention (VGD solution) according to the method of Tsushida, et al. [J. Jap. Soc. for Food Science and Technology,

41(9), 611 (1994)].

The test solutions were examined for the ability to inhibit autoxidation of linoleic acid based on the inhibition of fading of  $\beta$ -carotene color. In 10 ml of chloroform were respectively dissolved 10 mg of  $\beta$ -carotene, 1 g of linoleic acid and 2 g of Tween 80, and the solutions (0.5 ml, 0.2 ml and 1.0 ml, respectively) were put into a 200-ml Erlenmeyer flask. After the air in the system was substituted by nitrogen, the mixture in the flask was dried and then dissolved in 100 ml of distilled water. The obtained solution (45 ml) was mixed with 4 ml of 0.2 mol/liter phosphate buffer (pH 6.0) to prepare a substrate solution. The substrate solution (2.9 ml) was mixed with 0.1 ml of the cooking liquor of the present invention obtained in Example 4. The mixture was heated in a thermostat at 60°C and the change in absorbance (OD) at 470 nm was measured. Separately, a solution of 790 ppm VGD in acetonitrile and a commercial sake were respectively added to the substrate solution and the change in OD was measured in the same manner. The rate (%) of OD value measured 24 hours after the start of oxidation of linoleic acid based on the OD value at the start of oxidation is shown in Table 7. Oxidation of linoleic acid was inhibited by the addition of the cooking liquor of the present invention and VGD.

Table 7

	OD* (%)
Cooking liquor of the invention	27.3
Commercial sake	14.8
VGD (790 ppm)	36.7
Control	1.1

\*: Rate of OD value measured 24 hours after the start of oxidation of linoleic acid based on the OD value at the start of oxidation

#### Example 12

The *moromi* obtained in Example 4 was kept at 60°C in the dark for 14 days, whereby a cooking liquor of the present invention containing 25 ppb VGD was obtained. The 4VG

concentration in this cooking liquor was 0.01 ppm.

Example 13

To 100 ml of water were added 65 g of rice *koji*, 35 g of steamed rice, 0.1 g of yeast (Dia Yeast; trademark, Kyowa Hakko Kogyo Co., Ltd.) and 1 ml of lactic acid, followed by fermentation at 15°C for 2 days to obtain a yeast culture (first stage fermentation). To the resultant mixture were added 135 g of steamed rice, 65 g rice *koji*, 0.8 g of Sumizyme AC (trademark for hemicellulase produced by Shinnihon Kagaku Co., Ltd., 6400 units/g) and 200 ml of water, followed by initial reaction at 15°C for 2 days. Then, to the resultant mixture were added 235 g of steamed rice, 65 g rice *koji* and 400 ml of water, followed by intermediate reaction at 15°C for 2 days. Thereafter, to the resultant mixture were added 330 g of steamed rice, 70 g rice *koji* and 500 ml of water, followed by final reaction at 15°C for 19 days. The obtained fermentation product was filtered and the filtrate was pasteurized to obtain *moromi*. The *moromi* obtained above was kept in a thermostat with fluorescent illumination at 25°C for 24 hours to obtain sake. In the above example, the second stage fermentation is composed of intermittent reactions. There is no limit to the number of reactions. It takes three reactions in the above, for example as initial reaction, intermediate reaction and final reaction. It also be able to take two reaction, for example as initial reaction and final reaction.